

# Algebraic number theory

## 1. Introduction

Basic problem: given  $F \in \mathbb{Z}[X_1, \dots, X_n]$ , find  $x_1, \dots, x_n \in \mathbb{Z}$  s.t.  $F(x_1, \dots, x_n) = 0$ .

Equations of this form  $F(X_1, \dots, X_n) = 0$  are called Diophantine equations

Examples 1) Fermat's problem: for  $n \geq 3$

$\nexists$  non-trivial integral solutions to

$$X^n + Y^n = Z^n,$$

i.e. if  $(x, y, z) \in \mathbb{Z}^3$  is a solution, then  $xyz = 0$ .

Solved by Andrew Wiles (1995)

2) Sums of squares (Euler?). Let  $p$  be an odd prime. Can we find  $x, y \in \mathbb{Z}$  s.t.

$$p = x^2 + y^2.$$

In this example  $F = X^2 + Y^2 - p$

Answer: it is possible iff  $p \equiv 1 \pmod{4}$

Also possible for  $p = 2$ .

3) Pell's equation: let  $d > 1$  be a square-free integer (i.e. not divisible by  $p^2 \quad \forall \text{ prime } p$ ). Consider

$$X^2 - dY^2 = 1$$

This equation has  $\infty$  solutions.

## 2. Sums of squares

Fix odd prime  $p$ . Consider

$$X^2 + Y^2 = p \quad (*)$$

Rewrite:  $(X + iY)(X - iY) = p \quad i^2 = -1$

$x + iy = z$ , then  $x^2 + y^2 = p \Leftrightarrow z \cdot \bar{z} = p$

Def The ring of Gaussian integers

$$\mathbb{Z}[i] = \{x + iy \in \mathbb{C} \mid x, y \in \mathbb{Z}\}$$

Remark "Ring" will always mean a commutative ring with identity.

Solving  $(*) \Leftrightarrow$  finding  $z \in \mathbb{Z}[i]$ , s.t.  $z \cdot \bar{z} = p$

Def A ring  $R$  is

1) an integral domain, if  $\forall a, b \in R$

$$ab = 0 \Rightarrow a = 0 \text{ or } b = 0$$

2) a Euclidean domain, if it is integral  
s.t.  $\exists$  a map  $N: R \rightarrow \mathbb{Z}_{\geq 0}$  with

a)  $N(0) = 0$

b)  $\forall x, y \in R, y \neq 0 \quad \exists a, b \in R$  s.t.  
 $x = ay + b$  and  $N(b) < N(y)$

This means: we can <sup>or  $b=0$</sup>  divide  $x$  by  $y$   
with "remainder"  $b$ .

Example  $R = K[x]$   $K = \text{a field}$

$R$  is Euclidean dom. with  $N(f) = \deg(f)$   
for  $f \in K[x]$


Proposition 2.1  $\mathbb{Z}[i]$  is a Euclidean domain.

Proof  $\mathbb{Z}[i]$  integral, because it is a  
subring of  $\mathbb{C}$ . Define  $N(z) = z \cdot \bar{z} = |z|^2$

Let  $x, y \in \mathbb{Z}[i], y \neq 0$

Need:  $x = ay + b, \quad N(b) < N(y)$

$y$  and  $iy$  are linear indep.  $\mathbb{R}$  vectors  
in  $\mathbb{R}^2 \cong \mathbb{C}$



Elements of the form  $ay$ ,  
with  $a \in \mathbb{Z}[i]$  form a lattice

$$\mathbb{Z} \cdot y \oplus i\mathbb{Z}y$$

$$x = \alpha y + \beta iy, \quad \alpha, \beta \in \mathbb{R}$$

$$\text{Let } \alpha = \alpha_0 + \alpha_1 i, \quad \alpha_0 \in \mathbb{Z}, -\frac{1}{2} \leq \alpha_1 < \frac{1}{2}$$

$$\beta = \beta_0 + \beta_1 i, \quad \beta_0 \in \mathbb{Z}, -\frac{1}{2} \leq \beta_1 < \frac{1}{2}$$

$$x = \underbrace{(\alpha_0 + i\beta_0)}_a \cdot y + \underbrace{(\alpha_1 + i\beta_1)}_b y$$

$$b = x - a \cdot y \in \mathbb{Z}[i]$$

$$N(b) = |\alpha_1 + i\beta_1|^2 \cdot N(y) =$$

$$= (\alpha_1^2 + \beta_1^2) \cdot N(y) \leq \left(\frac{1}{4} + \frac{1}{4}\right) \cdot N(y)$$

$$= \frac{1}{2} N(y) < N(y). \quad \square$$

Recall: 1) an int. domain  $R$  is called  
principal ideal domain (PID) if  
any ideal in  $R$  is generated by  
one element.

2) Prop. 2.2  $R$  Euclidean domain  $\Rightarrow R$  PID

Idea of proof:  $I \subset R$  ideal.

Pick  $0 \neq x \in I$  s.t.

$$\forall 0 \neq y \in I \quad N(x) \subseteq N(y)$$

Then check that  $I = (x)$   $\square$

3) an element  $x \in R$  is a unit, if

$\exists y \in R$  s.t.  $xy = 1$ . Set of all units denoted by  $R^\times$

4) Let  $x \in R$  s.t.  $x \neq 0$ ,  $x \notin R^\times$   
 $x$  is called irreducible if  $x = ab$ ,  $a, b \in R$   
 $\Rightarrow$  either  $a \in R^\times$  or  $b \in R^\times$

otherwise  $x$  is called reducible

5) An int. domain  $R$  is called unique factorization domain, if  $\forall 0 \neq x \in R$   
s.t.  $x \notin R^\times$   $\exists$  a decomposition

$$x = p_1 \cdots p_n \quad \text{with } p_i \text{ are irred.}$$

and if  $x = q_1 \cdots q_m$  another decomposition with  $q_j$  irred, then

$n = m$  and after permuting  $q_j$

we have  $p_i = a_i q_i$  for some  $a_i \in R^\times$

6) Prop 2.3  $R$  is PID  $\Rightarrow R$  is UFD

Corollary:  $\mathbb{Z}[i]$  is UFD and PID

7) Prop 2.4 Assume that  $R$  is UFD

Then  $0 \neq x \in R$  is irreducible

$\Leftrightarrow (x)$  is prime, i.e.  $R/(x)$  is integral.

Prop 2.5  $z \in \mathbb{Z}[i]^{\times} \Leftrightarrow N(z) = 1$

$\Leftrightarrow z \in \{\pm 1, \pm i\}$

Proof:  $z \in \mathbb{Z}[i]^{\times} \Rightarrow \exists w$  s.t.  $zw = 1$   
 $\Rightarrow N(z) \cdot N(w) = 1 \Rightarrow N(z) = N(w) = 1$

$\Rightarrow z \in \{\pm 1, \pm i\}$   $\square$

Back to the equation

$$X^2 + Y^2 = p \quad (*)$$

$p = \text{odd prime}$

Prop. 2.6  $(*)$  has an integral solution  
iff  $p$  is reducible as an element  
of  $\mathbb{Z}[i]$ .

Proof: 1)  $(*)$  solvable  $\Rightarrow \exists z \in \mathbb{Z}[i]$  s.t.

$z \cdot \bar{z} = p$ . But  $N(z) = N(\bar{z}) = p > 1$ ,

so neither  $z$  nor  $\bar{z}$  is a unit

$\Rightarrow p$  is reducible.

2) Assume  $p$  is reducible

$$\Rightarrow \exists z, w \in \mathbb{Z}[i], \quad N(z) > 1, \quad N(w) > 1$$

$$p = z \cdot w$$

$$N(p) = p^2 = N(z) \cdot N(w)$$

$$\Rightarrow N(z) = N(w) = p$$

$$\Rightarrow p = z \cdot \bar{z} \Rightarrow (*) \text{ is solvable } \square$$

Thm 2.7 An odd prime  $p$  can be represented as the sum of two squares iff  $p \equiv 1 \pmod{4}$

Proof:  $(*)$  is solvable  $\xLeftrightarrow{\text{Prop. 2.6}}$   $p$  is reducible in  $\mathbb{Z}[i]$

Prop 2.4

$\Leftrightarrow (p)$  is not a prime ideal in  $\mathbb{Z}[i]$ .

$\Leftrightarrow \mathbb{Z}[i]/(p)$  has zero-divisors.

$$\text{Note: } \mathbb{Z}[i] = \frac{\mathbb{Z}[x]}{(x^2+1)}$$

$$\frac{\mathbb{Z}[i]}{(p)} \simeq \frac{\mathbb{Z}[x]}{(x^2+1, p)} \simeq \frac{\mathbb{Z}/(p)[x]}{(x^2+1)} = \frac{\mathbb{F}_p[x]}{(x^2+1)}$$

$\mathbb{F}_p$  = the field with  $p$  elements

$\frac{\mathbb{F}_p[x]}{(x^2+1)}$  has zero-div.  $\xLeftrightarrow{\text{Prop 2.4}}$   $x^2+1$  is reducible

$\Leftrightarrow X^2+1$  has a root in  $\mathbb{F}_p$ ,

i.e.  $\exists u \in \mathbb{F}_p$ , s.t.  $u^2 = -1$

Lemma  $-1 \in \mathbb{F}_p$  is a square  $\Leftrightarrow p \equiv 1 \pmod{4}$

Proof:  $\mathbb{F}_p^\times = \mathbb{F}_p \setminus \{0\}$  is a cyclic group (exercise)

Let  $\omega \in \mathbb{F}_p^\times$  is a generator

then 
$$\begin{array}{ccc} \mathbb{Z}/(p-1) & \xrightarrow{\sim} & \mathbb{F}_p^\times \\ a & \longmapsto & \omega^a \end{array}$$

$$\frac{p-1}{2} \longmapsto \omega^{\frac{p-1}{2}} = -1$$

$-1$  is a square  $\Leftrightarrow \exists a \in \mathbb{Z}$ , s.t.

$$(\omega^a)^2 = \omega^{\frac{p-1}{2}} \Leftrightarrow 2a \equiv \frac{p-1}{2} \pmod{p-1}$$

$$\Leftrightarrow \exists a, b \in \mathbb{Z} : 2a = \frac{p-1}{2} + b \cdot (p-1)$$

if  $p = 1 + 4n$  take  $a = n$ ,  $b = 0$

if  $p = 3 + 4n$  then  $\frac{p-1}{2} = 1 + 2n$

odd, but  $2a - b(p-1)$

is even  $\Rightarrow$  not solvable

This proves the lemma and the thm  $\square$

## Ideas arising from this proof

- need to study extensions of  $\mathbb{Z}$  such as  $\mathbb{Z}[i]$ ;  $\mathbb{Z}[i]$  is the ring of integers in the field  $\mathbb{Q}(i)$   
Need to study finite extensions of  $\mathbb{Q}$  (number fields)
- important to understand factorization into irred. factors in rings like  $\mathbb{Z}[i]$   
Unfortunately, not all rings of integers in number fields are UFD.

Example  $R = \mathbb{Z}[\sqrt{-5}] = \{a + \sqrt{-5}b \in \mathbb{C} \mid a, b \in \mathbb{Z}\}$

$$N(a + \sqrt{-5}b) = a^2 + 5b^2$$

$R$  is not UFD. Consider  $6 = 2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5})$   
 $2$  and  $3$  are irred, e.g. let  $2 = z \cdot w$

$$N(z) \cdot N(w) = 4 \Rightarrow N(z) = 2 - \text{cont.}$$

$\Rightarrow 2$  is irred, but  $1 \pm \sqrt{-5}$  is not divisible by  $2$  in  $R \Rightarrow$  contradicts uniqueness of factor. into irred. factors.

- Solution to this problem: consider prime ideals instead of irred. elements.

