

5. Algebraic integers

Def A number field is a finite ext. of $\mathbb{Q}$

$$\begin{array}{ccc} \mathbb{Q} & \subset & K \\ \cup & & \cup \end{array}$$

$$\mathbb{Z} \subset \mathcal{O}_K \leftarrow \text{integers in } K$$

What properties should $\mathcal{O}_K$ naturally have?

- 1) $\mathcal{O}_K$ should be a subring of K containing $\mathbb{Z}$.
- 2) If K_1, K_2 - number fields, $G: K_1 \hookrightarrow K_2$

$$\text{then } x \in \mathcal{O}_{K_1} \Leftrightarrow G(x) \in \mathcal{O}_{K_2}$$

$$\text{Equivalently, } \mathcal{O}_{K_1} = K_1 \cap \mathcal{O}_{K_2}$$

Note: 2) $\Rightarrow \mathcal{O}_K$ is stable under automorph.

of K , in part. if K is Galois / $\mathbb{Q}$

$$x \in \mathcal{O}_K, \text{ then } g(x) \in \mathcal{O}_K \quad \forall g \in \text{Gal}(K/\mathbb{Q})$$

Then x is a root of the polynom:

$$f = \prod_{g \in \text{Gal}(K/\mathbb{Q})} (X - g(x)) = X^n + a_1 X^{n-1} + \dots + a_n$$

a_i are symm. poly's in $g(x)$, $g \in \text{Gal}(K/\mathbb{Q})$

$$\Rightarrow a_i \in \mathbb{Q}, \quad 1) \Rightarrow a_i \in \mathcal{O}_K$$

$$2) \Rightarrow a_i \in \mathbb{Q} \cap \mathcal{O}_K = \mathbb{Z}$$

Def $x \in K$, where K is a number field, is called alg. integer if x is a root of some monic poly $f \in \mathbb{Z}[x]$

($f = a_0 x^d + \dots + a_1$ is monic if $a_0 = 1$)

Prop 5.1 For $x \in K$ the following are equivalent (TFAE):

- 1) x is an alg. integer (in the sense of the above def.)
- 2) the monic min. poly of x has integral coefficients.

Proof 2) $\Rightarrow$ 1) obvious

1) $\Rightarrow$ 2) Let x be a root of monic $f \in \mathbb{Z}[x]$, $h \in \mathbb{Q}[x]$ be the monic min poly of x .

Then $f = h \cdot z$ for some $z \in \mathbb{Q}[x]$
note: z is monic.

Assume $h \notin \mathbb{Z}[x]$. Then $\exists$ a prime p that appears in denominator of some

coeff. of h .

Define: n_1 = maximum exponent of p
appearing in denominators
of coeff of h

n_2 = — " — of z

then $n_1 \geq 1$ by assumption

$$p^{n_1+n_2} f = \underbrace{(p^{n_1} h)}_{\text{have no } p \text{ in denominators}} \cdot \underbrace{(p^{n_2} z)}$$

$\Rightarrow$ can reduce coeff. mod p

$$p^{n_1} h \pmod{p} = \bar{h} \in \mathbb{F}_p[x]$$

$$p^{n_2} z \pmod{p} = \bar{z} \in \mathbb{F}_p[x]$$

By the choice of n_1, n_2 $\bar{h} \neq 0, \bar{z} \neq 0$

$$p^{n_1+n_2} f \pmod{p} = 0$$

$\Rightarrow$ get $0 = \bar{h} \cdot \bar{z}$ — contradiction,

because $\mathbb{F}_p[x]$ has no zero
divisors. $\square$

We want to prove that σ_K is a
ring; first we generalize the notion
of an integer. Let $R \subset S$ be two
rings

Def 1) $x \in S$ is integral over R , if $\exists$ a monic $f \in R[x]$, s.t. $f(x) = 0$

2) $R \subset S$ is called integral ext., if $\forall x \in S$ is int / R

3) The integral closure of R in S
 $\bar{R} = \{x \in S \mid x \text{ is int. / } R\}$

Prop. 5.2 In the above setting TFAE

1) $x \in S$ is int. / R

2) x is contained in some subring $S' \subset S$, s.t. S' is a finitely generated R -module.

Rmk 2) means: $\exists s_1, \dots, s_n \in S'$, s.t.

$$R^{\oplus n} \longrightarrow S' \text{ is surj.}$$
$$(r_1, \dots, r_n) \longmapsto \sum_{i=1}^n r_i s_i$$

Proof of 5.2

1) $\Rightarrow$ 2) we have an equation of integral dependence

$$x^n + a_1 x^{n-1} + \dots + a_n = 0 \quad a_i \in R \quad (*)$$

Let S' be R -submodule of S spanned by $1, x, \dots, x^{n-1}$.

Then S' is a subring:

x^N can be expressed as a linear comb of $1, x, \dots, x^{n-1}$ using (*) for any $N \geq 0$

2) $\Rightarrow$ 1) Assume S' is generated by $s_1, \dots, s_n$; $x \in S' \Rightarrow$

$$(**) \quad x \cdot s_i = \sum_{j=1}^n a_{ij} s_j \quad \text{for some } a_{ij} \in R$$

Consider the matrix with elements in S'

$$A = \begin{pmatrix} a_{11} - x & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} - x & \dots & a_{2n} \\ \vdots & & & \\ a_{n1} & a_{n2} & \dots & a_{nn} - x \end{pmatrix}$$

$$\text{Note: } A \begin{pmatrix} s_1 \\ \vdots \\ s_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix} \quad \text{by } (**)$$

It follows (see Lemma below) that

$$\forall i \quad \det A \cdot s_i = 0$$

Since S' is a subring, we have

$$1 = \sum_{i=1}^n \alpha_i s_i \quad \text{for some } \alpha_i \in R$$

$$\Rightarrow \det A = \sum \alpha_i \underbrace{s_i \det A}_{0} = 0$$

$\det A = (-1)^n x^n + b_1 x^{n-1} + \dots + b_n \quad b_i \in R$
 multiplying by $(-1)^n \Rightarrow$ get equation
 of int. depend. for x . $\square$

Lemma R arbitrary ring, $A = (a_{ij})$
 a $n \times n$ matrix $a_{ij} \in R$, $s_1, \dots, s_n \in R$
 Assume $A \begin{pmatrix} s_1 \\ \vdots \\ s_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}$. Then

$$\forall i \quad \det A \cdot s_i = 0$$

Proof Let $A^+ = (A_{ij})$, where
 $A_{ij} = (-1)^{i+j} \cdot \det(\underbrace{(j,i)\text{-th minor of } A}_{(n-1) \times (n-1) \text{ matrix obtained from } A \text{ by removing } j\text{-th row and } i\text{-th column}})$

Then:

$$A^+ \cdot A = \det A \cdot Id$$

$$\Rightarrow \underbrace{0}_{0} = A^+ \cdot A \cdot \begin{pmatrix} s_1 \\ \vdots \\ s_n \end{pmatrix} = \begin{pmatrix} \det A \cdot s_1 \\ \vdots \\ \det A \cdot s_n \end{pmatrix} \quad \square$$

Prop. 5.3 1) $R \subset S$ ring extension

$\Rightarrow$ the int. closure $\bar{R}$ is a subring of S , containing R

2) $R \subset S \subset T$ rings, assume $R \subset S$ is an integral ext.

If $x \in T$ is int / $S \Rightarrow x$ is int / R

(transitivity of integrality)

Proof 1) Need: $x_1, x_2 \in \bar{R} \Rightarrow x_1 + x_2 \in \bar{R}$
 $x_1, x_2 \in \bar{R}$

$$(x) \begin{cases} x_1^n + a_1 x_1^{n-1} + \dots + a_n = 0 & a_i \in R \\ x_2^m + b_1 x_2^{m-1} + \dots + b_m = 0 & b_i \in R \end{cases}$$

We can assume $n > 1, m > 1$
(case $n=1$ or $m=1$ — exercise)

Let S' be R -submodule of S generated by $x_1^i x_2^j$ for $i=0 \dots n-1$
 $j=0 \dots m-1$

S' is lin. generated

Since any $x_1^u x_2^v, u, v \geq 0$ can be expressed as a lin. comb. of

$$x_1^i x_2^j \quad i=0 \dots n-1, j=0 \dots m-1$$

using (*). Hence S' is a subring.

By construction S' contains $x_1 + x_2$
and $x_1 x_2 \xRightarrow{\text{Prop. 5.2}} x_1 + x_2, x_1 x_2 \in \overline{R} \cap$

$$2) x \in T \text{ int} / S \Rightarrow x^n + s_1 x^{n-1} + \dots + s_n = 0$$

For some $s_i \in S$; s_i are int / R

$\Rightarrow$ the subring of S generated by

$s_1, \dots, s_n$ is spanned / R by a finite number of monomials

$$s_1^{k_1} \dots s_n^{k_n}, \quad 0 \leq k_i < N$$

Consider $T' =$ subring of T

spanned by $x^i s_1^{k_1} \dots s_n^{k_n}, \quad 0 \leq i < n$
 $0 \leq k_i < N$

T' is fin. gen $\xRightarrow{\text{Prop 5.2}} x$ is int / R $\square$

Def For a ring ext $R \subset S$

R is int. closed in S if $\overline{R} = R$

Cor. from Prop 5.3 For $R \subset S$

then $\overline{R}$ is int. closed in S

$$\text{i.e. } \overline{\overline{R}} = \overline{R}$$

Proof Apply Prop 5.3 (2) to
 $R \subset \overline{R} \subset S$ $\square$

Back to the number fields

$$\mathbb{Z} \subset \mathbb{Q} \subset K$$

$\Downarrow$

$\mathbb{Z} \subset K$ a ring extension;

By definition $\mathcal{O}_K = \text{int. closure}$
of $\mathbb{Z}$ in K

In part. Prop 5.3 $\Rightarrow \mathcal{O}_K$ is a ring,
int. closed in K

Example Let $K = \mathbb{Q}(i)$

Min poly of $z = a + ib \in \mathbb{Q}(i)$

$$X^2 - (\sigma_1(z) + \sigma_2(z))X + \sigma_1(z) \cdot \sigma_2(z) \\ = X^2 - 2aX + a^2 + b^2$$

$$z \in \mathcal{O}_K \Leftrightarrow \begin{cases} 2a \in \mathbb{Z} \\ a^2 + b^2 \in \mathbb{Z} \end{cases} \Leftrightarrow \begin{cases} a = \frac{n}{2}, n \in \mathbb{Z} \\ \frac{n^2}{4} + b^2 = m \in \mathbb{Z} \end{cases}$$

$$\frac{n^2}{4} + b^2 = m \Rightarrow n^2 = 4m - 4b^2$$

$$n \equiv 0 \pmod{2} \Rightarrow a \in \mathbb{Z} \Rightarrow b^2 \in \mathbb{Z} \\ \Rightarrow b \in \mathbb{Z}$$

$$n \equiv 1 \pmod{2} \Rightarrow n^2 \equiv 1 \pmod{4}$$

$$\text{from } n^2 = 4m - 4b^2$$

$$4b^2 \stackrel{\downarrow}{\in} \mathbb{Z} \Rightarrow b = \frac{l}{2} \quad l \in \mathbb{Z}$$

$$m = \frac{n^2}{4} + b^2 = \frac{n^2 + l^2}{4} \in \mathbb{Z}$$

l^2 is either 0 or 1
mod 4

$$\Rightarrow n^2 + l^2 \equiv 1 \pmod{4}$$

$$\text{or } n^2 + l^2 \equiv 2 \pmod{4}$$

in any case $\frac{n^2 + l^2}{4} \notin \mathbb{Z}$

$\Rightarrow n \equiv 1 \pmod{2}$ can not happen.

$$\Rightarrow \mathcal{O}_K \simeq \mathbb{Z}[i]$$