

Exercises, Algebraic Geometry II – Week 13

Exercise 56. (2 points) *Projection formula.*

Let $f: X \rightarrow Y$ be a morphism and $\mathcal{F} \in \text{Qcoh}(X)$. Show that for a locally free $\mathcal{G} \in \text{Coh}(Y)$ one has

$$R^i f_*(\mathcal{F} \otimes f^* \mathcal{G}) \cong R^i f_*(\mathcal{F}) \otimes \mathcal{G}.$$

Exercise 57. (4 points) *Künneth formula.*

Let X_1 and X_2 be separated schemes over a field k . One writes $\mathcal{F}_1 \boxtimes \mathcal{F}_2$ for $p_1^* \mathcal{F}_1 \otimes p_2^* \mathcal{F}_2$ for quasi-coherent sheaves \mathcal{F}_1 on X_1 and \mathcal{F}_2 on X_2 .

1. Show that

$$H^n(X_1 \times_k X_2, \mathcal{F}_1 \boxtimes \mathcal{F}_2) \cong \bigoplus_{i+j=n} H^i(X_1, \mathcal{F}_1) \otimes_k H^j(X_2, \mathcal{F}_2).$$

2. Prove that $R^n p_{1*}(\mathcal{F}_1 \boxtimes \mathcal{F}_2) \cong \mathcal{F}_1 \otimes_k H^n(X_2, \mathcal{F}_2)$. Compare this with the assumptions on the previous exercise.

Exercise 58. (2 points) *Hodge numbers of products.*

For a smooth projective variety X over a field k one defines $h^{p,q}(X) := \dim_k H^q(X, \Omega_{X/k}^p)$. Compute the Hodge numbers $h^{p,q}(X_1 \times_k X_2)$ of a product of two smooth projective varieties X_1, X_2 .

Exercise 59. (3 points) *Hodge bundles.*

Let $f: X \rightarrow Y$ be a smooth projective morphism with Y Noetherian and $\dim X = \dim Y + 1$. Show that the Hodge bundles $R^q f_*(\Omega_{X/Y}^p)$ are locally free sheaves. (In characteristic zero, this holds true without the assumption on the dimension.)

Exercise 60. (3 points)

Let $f: X \rightarrow Y$ be a projective morphism with Y Noetherian integral of positive dimension. Let $\mathcal{F} \in \text{Coh}(X)$ be flat over Y and such that there exists at most one $y \in Y$ with $H^0(X_y, \mathcal{F}_y) \neq 0$. Show that then $f_* \mathcal{F} = 0$. (Convince yourself that the flatness of \mathcal{F} is really needed for this.)

Exercise 61. (2 points)

Let Y be a Noetherian scheme and \mathcal{E} a locally free sheaf of rank $n + 1$ on Y . Let $\pi: X = \mathbb{P}(\mathcal{E}) \rightarrow Y$ and $\mathcal{O}(1)$ be the corresponding invertible sheaf on X . Denote by $\omega_{X/Y}$ the relative canonical sheaf. Using the isomorphism $R^n \pi_* \omega_{X/Y} \simeq \mathcal{O}_Y$ (which one can deduce from Serre duality) and the relative Euler sequence (see exercises 11 and 34) prove that for any $l \in \mathbb{Z}$ we have $R^n \pi_*(\mathcal{O}(l)) \simeq \pi_*(\mathcal{O}(-l - n - 1))^\vee \otimes (\Lambda^{n+1} \mathcal{E})^\vee$.

Due Monday 18 July, 2016.

Attention: no lecture on Thursday 14.07.2016. Next lecture on Monday 18.07.2016

Exams:

- 02.08.2016, 9.00 - 11.00, Großer Hörsaal, Wegelerstr. 10;
- 28.09.2016, 9.00 - 11.00, Großer Hörsaal, Wegelerstr. 10.